


# Study of $\Upsilon(1S)$ flow in pPb collision system with the CMS detector

**KiSoo Lee**


September 3rd, 2022



# Particle anisotropy





- Due to the length and pressure difference, spatial anisotropy converted into a momentum anisotropy
- Non-zero  $v_2$  is observed even in the small system




PLB 765 (2017) 193

# Quarkonia flow


- Quarkonia are expected to carry out information of the initial state and the medium effects
- Large  $v_2$  of  $J/\psi$  at low- $p_T$  from recombination effect while  $v_2$  is zero for  $\Upsilon(1S)$  in PbPb
- Non-zero  $v_2$   $J/\psi$  is observed in pPb
- $\Upsilon(1S)$   $v_2$  in pPb is not measured yet



CMS-PAS-HIN-21-008




PLB 819 (2021) 136385



PLB 791 (2019) 172

# Reconstruction




# Same event correlation

$$\Delta\eta = \eta^\Upsilon - \eta^{\text{trk}}$$

$$\Delta\phi = \phi^\Upsilon - \phi^{\text{trk}}$$

$$S(\Delta\eta, \Delta\phi) = \frac{1}{N_{\text{trig}}} \frac{d^2 N^{\text{same}}}{d\Delta\eta d\Delta\phi}$$

←  $\Upsilon$   
← charged track



- Two-particle correlations in  $\Delta\eta$ - $\Delta\phi$  ( $\Upsilon$ -track)
- $\Upsilon$ : trigger, track: associator
- $0.3 < p_{\text{T}}^{\text{track}} < 3$


# Mixed event correlation

$$\Delta\eta = \eta^\gamma - \eta^{\text{trk}}$$

$$\Delta\phi = \phi^\gamma - \phi^{\text{trk}}$$

$$B(\Delta\eta, \Delta\phi) = \frac{1}{N_{\text{trig}}} \frac{d^2 N^{\text{mix}}}{d\Delta\eta d\Delta\phi}$$

←  $\gamma$   
← charged track



- The  $\gamma$  candidate as trigger particle correlated with the charged track associators from the different event
- 10 random event mixed within  $|z_{\text{vtx}}^1 - z_{\text{vtx}}^2| < 2 \text{ cm}$







# Two-particle correlation method

$$S(\Delta\eta, \Delta\phi) = \frac{1}{N_{trig}} \frac{d^2 N^{same}}{d\Delta\eta d\Delta\phi}$$

$$B(\Delta\eta, \Delta\phi) = \frac{1}{N_{trig}} \frac{d^2 N^{mix}}{d\Delta\eta d\Delta\phi}$$

$$\frac{1}{N_{trig}} \frac{d^2 N^{pair}}{d\Delta\eta d\Delta\phi} = B(0,0) \times \frac{S(\Delta\eta, \Delta\phi)}{B(\Delta\eta, \Delta\phi)}$$





PLB 765 (2017) 193

- Cancel out the random combinatorial background and acceptance effects

# Observed $V_2$ extraction

PLB 765 (2017) 193




- Long-range ( $|\Delta\eta| > 1$ ) events projected to  $\Delta\phi$  axis in order to reject jet contribution
- $V_n(\Upsilon\text{-trk})$  is determined from a Fourier decomposition

$$\frac{1}{N_{\text{trig}}} \frac{d^2 N^{\text{pair}}}{d\Delta\eta d\Delta\phi} = \frac{N_{\text{assoc}}}{2\pi} \left\{ 1 + \sum_n 2V_{n\Delta} \cos(n\Delta\phi) \right\}$$

$V_n$ :  $\Upsilon$ -track  
 $v_n$ :  $\Upsilon$



# Simultaneous fitting




- Observed  $V_2$  is composed of signal-track correlation and background-track correlation
- To extract signal  $V_2$ , simultaneous fitting with mass is applied

# Non-flow subtractions

PLB 765 (2017) 193

- Low-multiplicity subtraction to remove non-flow effect (mostly from back-to-back jet correlation)
- Jet yield ratio used to account for the enhanced jet correlations from low to high-multiplicity

CMS pp  $\sqrt{s} = 13$  TeV



$$V_2^{\text{sub}} = V_2^{\text{Sig}}(\text{high}) - V_2^{\text{Sig}}(\text{low}) \times \frac{N_{\text{assoc}}(\text{low})}{N_{\text{assoc}}(\text{high})} \times \frac{J_{\text{jet}}(\text{high})}{J_{\text{jet}}(\text{low})}$$




# Non-flow subtractions

PLB 765 (2017) 193

- Low-multiplicity subtraction to remove non-flow effect (mostly from back-to-back jet correlation)
- Jet yield ratio used to account for the enhanced jet correlations from low to high-multiplicity

CMS pp  $\sqrt{s} = 13$  TeV




$$V_2^{\text{sub}} = V_2^{\text{Sig}}(\text{high}) - V_2^{\text{Sig}}(\text{low}) \times \frac{N_{\text{assoc}}(\text{low})}{N_{\text{assoc}}(\text{high})} \times \frac{J_{\text{jet}}(\text{high})}{J_{\text{jet}}(\text{low})}$$

# Track $V_2$ subtractions

$$\Delta\eta = \eta^{\text{trk}} - \eta^{\text{trk}}$$

$$\Delta\phi = \phi^{\text{trk}} - \phi^{\text{trk}}$$





$$v_2^{\text{sub}} = \frac{V_2^{\text{sub}}}{\sqrt{V_2^{\text{sub}}(\text{trk})}}$$

- To extract pure  $\Upsilon(1S)$   $v_2$ , track  $v_2$  is divided from the  $\Upsilon(1S)$ -track  $v_2$





# Result



- Y(1S)  $v_2$  is consistent with 0 regardless of the system size
- Hint of different behavior for charmonia and bottomonia

# Summary

- $v_2$  of quarkonia is useful tool to study the path-length dependent modification effect and collectivity of heavy flavors
- $\Upsilon(1S)$   $v_2$  measured for the first time in pPb
- $\Upsilon(1S)$   $v_2$  is close to 0 regardless of the system size
- Hint of different behavior for charmonia and bottomonia
- Presented in QM, SQM, Target journal: PLB



**Back up**

# CMS detector

Calorimeters  
(Electromagnetic & Hadron)

